

Verbal Rating Scale (VRS)

Verscheidene auteurs

Meetinstrument	Verbal Rating Scale
Afkorting	VRS
Auteur	Verscheidene auteurs
Thema	Symptoommanagement pijn
Doel	Pijn beoordelen bij de patiënt
Populatie	Niet gespecificeerd
Afname	Zorgverlener
Aantal items	1 item
Aanwezigheid patiënt vereist	Ja
Vindplaats meetinstrument	Herr, K. A. & Mobily, P. R. (1993). Comparison of selected pain assessment tools for use with the elderly. <i>Appl.Nurs.Res.</i> , 6, 39-46.

Doel

Pijncontrole bij de patiënt aan de hand van zelfrapportage op een één-item schaal met verschillende pijndescriptoren.

Doelgroep

De Verbal Rating Scale (VRS) werd reeds in een brede populatie gevalideerd; dit zowel bij volwassenen als bij ouderen (waaronder ook ouderen met een verminderd cognitief functioneren). Studies die de betrouwbaarheid en validiteit van de VRS onderzoeken bij kinderen, zijn echter schaars.

Onderzoek naar de validiteit van de VRS wordt o.a. gerapporteerd bij chirurgische patiënten (Jensen, Chen & Brugger, 2002), ouderen (Herr, Spratt, Mobily & Richardson, 2004), CVA-patiënten (Benaim et al., 2007); traumapatiënten (Puntillo & Neighbor, 1997), dementerenden (Chibnall & Tait, 2001; Kaasalainen & Crook, 2003; Pautex et al., 2005; Pautex et al., 2006; Taylor & Herr, 2003; Ware, Epps, Herr & Packard, 2006), oncologiepatiënten (Paice & Cohen, 1997), kinderen (Wong & Baker, 1988),....

Beschrijving

De VRS als meetinstrument omvat verschillende pijndescriptoren die gerangschikt zijn volgens toenemende pijnintensiteit. Er bestaan echter tal van varianten op de VRS. Veelgebruikte pijndescriptoren zijn o.a. 'geen pijn – zwak – matig – hevig – extreem' en 'geen pijn – zeer zwak – mild – lastig – hevig – ondraaglijk'.

Laatstgenoemde combinatie is een onderdeel van de McGill Pain Questionnaire, de Present Pain Inventory (PPI), en wordt vaak als VRS gehanteerd (Melzack, 1975). De woorden worden gescoord door aan elk adjetief een getal toe te kennen (0 – 5).

Andere benamingen voor de VRS zijn Verbal Descriptor Scale (VDS), Verbal Pain Scale (VPS) en Simple Descriptor Scale (SDS). Een soortgelijke variant is ook de Iowa Pain Thermometer (Herr, Spratt, Garand & Li, 2007). Hierbij gaat het telkens om gelijkaardige meetinstrumenten. Kleine verschillen kunnen terug te vinden zijn in de adjetieven die men gebruikt tussen de verschillende meetinstrumenten.

Betrouwbaarheid

De betrouwbaarheid van de VRS werd uitvoerig bestudeerd in de studies van Benaim et al. (2007), Chibnall & Tait (2001), Herr et al. (2004), Kaasalainen & Crook (2003), Pautex et al. (2005), Pautex et al. (2006), Taylor & Herr (2003), Ware et al. (2006), Wong & Baker (1988).

Met uitzondering van de studie van Chibnall & Tait (2001) en Benaim et al. (2007) levert de *test – hertest* vrij sterke correlaties op ($r > 0.70$). Deze correlaties liggen lager voor ouderen met een cognitief verminderd functioneren, hoewel de range nog steeds 0.40 – 0.97 bedraagt. Kaasalainen & Crook (2003) concluderen dan ook dat de VRS een betrouwbaar en adequaat meetinstrument is voor dementerende personen.

Cronbach's alpha tussen de VRS en andere één-item meetschalen (VAS – Numeric Rating Scale – Verbal Numeric Scale – Faces Pain Scale) bedroeg $\alpha = 0.97$ (Herr et al., 2004).

De *interrater reliability* leverde een correlatie op van respectievelijk $r = 0.46 – 0.52$, $r = 1.00$, $r = 0.80$ en $r = 0.95$ in de studies van Benaim et al. (2007), Herr et al. (2004) Pautex et al. (2005) en Pautex et al. (2006).

Validiteit

De VRS is een valide meetinstrument in het controleren van pijn bij de patiënt. De *concurrent validity* werd nagegaan in de studies van Benaim et al. (2007), Jensen, Engel, McKearnan & Hoffman (2003), Paice & Cohen (1997), Pautex et al. (2005), Puntillo & Neighbor (1997), Taylor & Herr (2003) en Ware et al. (2006). De correlatie tussen de VRS en ander pijnschalen is overwegend sterk; in het merendeel van de studies wordt een correlatie hoger dan 0.70 gerapporteerd.

Om de *construct validity* van de VRS na te gaan, hebben Herr et al. (2004) en Jensen et al. (2003) een *principale componenten analyse* uitgevoerd. Uit de pijnscores op de verschillende één-item meetinstrumenten werd 1 factor geabstraheerd. Een *factor-analyse* op de verschillende VRS-pijnscores uit herhaalde metingen gedurende 12 dagen, resulteerde eveneens in 1 factor (Chibnall & Tait, 2001).

De *convergent validity* werd bestudeerd in het onderzoek van Jensen et al. (2003) en Kaasalainen & Crook (2003) door de VRS te correleren aan respectievelijk

constructen zoals depressie en gedragsobservaties bij pijn. De gerapporteerde correlaties waren matig. Dit kan er op wijzen dat deze gerelateerde constructen zich enigszins onderscheiden van pijn en de VRS enkel het construct 'pijn' meet.

De VRS is eveneens sensitief aan pijnprikkels met toenemende pijnintensiteit (Herr et al., 2004) en het behandelen van pijn d.m.v. analgetica (Jensen et al., 2002).

Gebruiksvriendelijkheid

De gebruiksvriendelijkheid van dit instrument werd niet bestudeerd. Rekening houdend met de omvang van het meetinstrument, kan men er van uitgaan dat de VRS vlot en snel kan afgenummeren worden. Wanneer gepeild wordt naar de voorkeur van patiënten voor een bepaald meetinstrument, scoort de VRS behoorlijk goed (Herr et al., 2004; Paice & Cohen, 1997; Puntillo & Neighbor, 1997). Tevens blijkt dat de VRS goed begrijpbaar is voor patiënten en weinig of geen problemen genereert bij afname (Herr et al., 2004; Paice & Cohen, 1997; Pautex et al. 2006; Taylor & Herr, 2003).

Opmerkingen

Een vergelijking tussen verschillende meetinstrumenten toont aan dat de VRS aangewezen kan zijn in het meten van pijn bij ouderen, en meer specifiek bij ouderen met een verhoogde leeftijd en dementerenden (Herr et al., 2004; Herr et al., 2007; Kaasalainen & Crook, 2003; Pautex et al., 2005; Pautex et al., 2006; Taylor & Herr, 2003; Ware et al., 2006). Op basis van hun resultaten, adviseren Kaasalainen & Crook (2003) en Pautex et al. (2005) de VRS bij dementerenden. Naast de VRS blijkt ook de FPS een betrouwbaar en valide instrument te zijn bij ouderen en personen met een verminderd cognitief functioneren (Freeman, Smyth, Dallam & Jackson, 2001; Pautex et al., 2006, Taylor & Herr, 2003, Ware et al., 2006).

Referenties

- Benaim, C., Froger, J., Cazottes, C., Gueben, D., Porte, M., Desnuelle, C., & Pelissier, J. Y. (2007). Use of the Faces Pain Scale by left and right hemispheric stroke patients. *Pain*, 128, 52-58.
- Chibnall, J. T. & Tait, R. C. (2001). Pain assessment in cognitively impaired and unimpaired older adults: a comparison of four scales. *Pain*, 92, 173-186.
- Freeman, K., Smyth, C., Dallam, L., & Jackson, B. (2001). Pain measurement scales: a comparison of the visual analogue and faces rating scales in measuring pressure ulcer pain. *J.Wound.Ostomy.Continence.Nurs.*, 28, 290-296.
- Herr, K. A., Spratt, K., Garand, L., & Li, L. (2007). Evaluation of the Iowa Pain Thermometer and Other Selected Pain Intensity Scales in Younger and Older Adult Cohorts Using Controlled Clinical Pain: A Preliminary Study. *Pain Medicine*, 6, 585-600.
- Herr, K. A., Spratt, K., Mobily, P. R., & Richardson, G. (2004). Pain intensity assessment in older adults: use of experimental pain to compare psychometric

properties and usability of selected pain scales with younger adults. *Clin.J.Pain*, 20, 207-219.

Jensen, M. P., Chen, C., & Brugger, A. M. (2002). Postsurgical pain outcome assessment. *Pain*, 99, 101-109.

Jensen, M. P., Engel, J. M., McKearnan, K. A., & Hoffman, A. J. (2003). Validity of pain intensity assessment in persons with cerebral palsy: a comparison of six scales. *J Pain*, 4, 56-63.

Kaasalainen, S. & Crook, J. (2003). A comparison of pain-assessment tools for use with elderly long-term-care residents. *Can J Nurs Res*, 35, 58-71.

Melzack, R. (1975). The McGill Pain Questionnaire: major properties and scoring methods. *Pain*, 1, 277-299.

Paice, J. A. & Cohen, F. L. (1997). Validity of a verbally administered numeric rating scale to measure cancer pain intensity. *Cancer Nurs*, 20, 88-93.

Pautex, S., Herrmann, F., Le Lous, P., Fabjan, M., Michel, J. P., & Gold, G. (2005). Feasibility and reliability of four pain self-assessment scales and correlation with an observational rating scale in hospitalized elderly demented patients. *J Gerontol.A Biol.Sci.Med.Sci.*, 60, 524-529.

Pautex, S., Michon, A., Guedira, M., Emond, H., Le Lous, P., Samaras, D., Michel, J. P., Herrmann, F., Giannakopoulos, P., & Gold, G. (2006). Pain in severe dementia: self-assessment or observational scales? *J Am Geriatr Soc*, 54, 1040-1045.

Puntillo, K. A. & Neighbor, M. L. (1997). Two methods of assessing pain intensity in English-speaking and Spanish-speaking emergency department patients. *J Emerg Nurs*, 23, 597-601.

Taylor, L. J. & Herr, K. (2003). Pain intensity assessment: a comparison of selected pain intensity scales for use in cognitively intact and cognitively impaired African American older adults. *Pain Manag.Nurs*, 4, 87-95.

Ware, L. J., Epps, C. D., Herr, K., & Packard, A. (2006). Evaluation of the Revised Faces Pain Scale, Verbal Descriptor Scale, Numeric Rating Scale, and Iowa Pain Thermometer in older minority adults. *Pain Manag.Nurs*, 7, 117-125.

Wong, D. L. & Baker, C. M. (1988). Pain in children: comparison of assessment scales. *Pediatr Nurs*, 14, 9-17.

Zelman, D. C., Gore, M., Dukes, E., Tai, K. S., & Brandenburg, N. (2005). Validation of a modified version of the brief pain inventory for painful diabetic peripheral neuropathy. *J Pain Symptom Manage*, 29, 401-410.

Vindplaats meetinstrument

Herr, K. A. & Mobily, P. R. (1993). Comparison of selected pain assessment tools for use with the elderly. *Appl.Nurs.Res.*, 6, 39-46.

VERBAL RATING SCALE (VRS)

VERSCHEIDENE AUTEURS

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Jensen, M. P., Chen, C., & Brugger, A. M. (2002)	Not specified.	123 patients who had undergone knee surgery and 124 women who had undergone a laparotomy . (n = 227)	Repeated measures design: Baseline measurement with 16 additional measures up to 24h following surgery.		Sen
Chibnall, J. T. & Tait, R. C. (2001)	A proprietary subacute care facility.	Cognitively impaired and unimpaired older adults over 55 years. (n = 90)	Repeated measures design: Patients made ratings of current pain three times/day for 7 days. They also made retrospective daily, weekly, and bi-weekly ratings of usual, worst, and least pain levels over a 14-day period. Ratings were made on four different scales: a five-point verbal rating scale, a seven-point faces pain scale, a horizontal 21-point (0 ± 100) box scale, and a vertical 21-point (0 ± 20) box scales (measuring pain intensity).	S	CsV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
	<p>(Sen) <i>Repeated measures ANOVA:</i> Dependent variables: VAS and VRS pre- to posttreatment difference scores Independent variables: time and analgeticum treatment - VAS tended to be more sensitive than VRS showing the smallest F values ($p < 0.001$). - A composite measure made up of a standardized average of three measures of outcome (VAS difference score, VRS difference score and a pain relief rating was not more sensitive to treatment effects (analgetica) than any one measure.</p>	
<p>(S) <i>Intra-class correlation:</i> VRS' reliability coefficients for cognitively unimpaired and impaired patients was respectively $0.30 \leq r < 0.55$ and $0.15 < r < 0.40$.</p>	<p>(CsV) <i>Factor analyse:</i> For each day, 12 day-retrospective ratings (usual, worst, least pain for each of four scales) were subjected to a principal-axis factor analysis. A single 'pain intensity' factor emerged and factor loadings for VRS ranged from 0.55 to 0.80.</p>	The horizontal 21-point box scale emerged as the best scale with respect to both psychometrics and validity, regardless of mental status.

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Wong, D. L. & Baker, C. M. (1988)	Pediatric units of two general hospitals in the South-Central United States.	Hospitalized children in three age groups: 3 to 7 (n = 52), 8 to 12 (n = 52), and 13 to 18 years (n = 46). (n = 150)	Comparative study. The following six scales were compared on reliability and validity: Simple Descriptive Scale (SDS), NRS, FPS, the Glasses Scale, the Chips Scale, and the Color Scale.	S	CrV
Puntillo, K. A. & Neighbor, M. L. (1997)	A level 1 trauma center emergency department in San Francisco.	Ninety-five English-speaking patients (67 male and 28 female) and 21 Spanish-speaking patients (16 male and 5 female) participated in the study. Study patients were treated in the emergency department for sprains or strains (35%), fractures (19%), contusions (14%), cellulitis or abscesses (12%), or other miscellaneous conditions such as headaches or abdominal/flank pain (20%). (n = 59)	Repeated measures design: patients were asked to use the NRS and VRS (English or Spanish version according the language of the patient) seven times over a 2-hour period, once immediately before and six times after receiving an analgesic.		CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Test-retest reliability: Retest occurred the day after the initial test except in a few instances when it was done 2 to 4 days later. Percentage agreement for the 6 scales was:</p> <ul style="list-style-type: none"> - SDS: 72.73% - NRS: 75.44% - FPS: 74.24% - Glasses: 75.38% - Chips: 77.27% - Colors: 68.18% 	<p>(CrV) Concurrent validity: Children were asked to list painful events that they had experienced since being hospitalized and rank these from most to least painful. Each scale was also used to rate these painful events. A percentage consistency between the ranking of the events and the responses for each pain scale was calculated. Percentage agreement for the 6 scales was:</p> <ul style="list-style-type: none"> - SDS: 62.81% - NRS: 60.00% - FPS: 60.43% - Glasses: 63.70% - Chips: 69.06% - Colors: 58.39% 	<p>The FPS was the most preferred scale by all age groups.</p> <p>The finding of an increase in validity and reliability with age is consistent with children's advancing cognitive ability. However, reliability increased only from the 3 to 7 year age group. Reliability decreased in the 13 to 18 year age group for all the scales except for the color scale and the SDS.</p> <p>No one scale demonstrates superiority in validity or reliability. No significant differences exist among the scales for any age group.</p>
	<p>(CrV) Concurrent validity: Spearman correlations between scores on the two scales were computed for each of the seven data collection times. Correlations between the two scales were moderate to very high ($r = 0.48$ to 0.96) and statistically significant ($p < 0.05$ to $p < 0.001$) at each of the seven time periods. In fact, there was an increase in the relationship between the two pain scores (NRS and VRS) from the beginning to the end of the study.</p>	<p>Slightly more English speaking patients preferred the NRS (50) over the VRS (35), and slightly more Spanish-speaking patients preferred the VRS (11) over the NRS (9). However, these findings were not statistically significant.</p>

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Ware, L. J., Epps, C. D., Herr, K., & Packard, A. (2006)	Not specified.	68 subjects aged 60 years and older with cognitive impairments. The mean score for the Mini Mental State Examination (MMSE) was 23 (standard deviation = 5.4) with a range from 10 to 30. Fifty-nine percent (n = 40) of the sample scored 24 or greater indicating no cognitive impairment (CI). Forty-one percent (n = 28) scored less than 24 indicating some degree of CI. (n = 68)	Comparative study: Subjects were instructed to recall a vividly remembered pain and rate this remembered pain using the Iowa Pain Thermometer (IPT), the Verbal Descriptor Scale (VDS), a 0 to 10 Numeric Rating Scale (NRS), and the Faces Pain Scale Revised (FPS-R).	S	CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Test-retest reliability: The researcher returned 2 weeks later and reminded subjects of the “vividly remembered pain” identified at the first assessment and asked them to rate that pain again using all four scales. In the cognitively intact group, Spearman rank correlation coefficients between the two-week vividly remembered pain ratings were 0.87 (NRS), followed by the VDS (0.86), IPT (0.81), and FPS-R (0.76). Correlations ranged from 0.77 to 0.89 in the CI group. The FPS-R had the strongest test-retest reliability coefficient (0.89) in this group followed by the IPT (0.82), VDS (0.79), and NRS (0.77).</p>	<p>(CrV) Concurrent validity: Spearman correlations between pain ratings on the selected scales ranged from 0.64 to 0.90 in the cognitively intact group and from 0.56 to 0.83 in the CI group. The lowest correlations in the CI group and intact group (0.64–0.84) were found between the FPS-R and the other scales (0.56–0.66).</p>	<p>In terms of the concurrent validity, moderate to high inter-tool correlations for the CI and cognitively intact groups were found with the exception of low correlations associated with the FPS-R, suggesting that the FPS-R may measure overall affect as opposed to pain.</p> <p>Four participants with moderate CI were unable to follow directions and complete the VDS and IPT. The NRS had the highest failure rate with six participants with moderate CI and one mildly impaired participant unable to use the scale. No failures occurred when using the FPS-R to evaluate pain.</p> <p>The NRS ($n = 12/36, 33\%$) was the preferred scale in the cognitively intact group, and the FPS-R ($n = 13/24, 54\%$) was the preferred scale in the CI group.</p>

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Jensen, M. P., Engel, J. M., McKearnan, K. A., & Hoffman, A. J. (2003)	Not specified.	Persons with cerebral palsy (CP) who have reported the presence of a chronic pain problem. Pain intensity assessment data were available for 24 participants from the survey study and 45 participants from the longitudinal study (n = 69)	Comparative study . Study participants were recruited from 2 other ongoing studies (a single-assessment survey and a 2-year longitudinal study). A subgroup of those who reported ongoing problems with pain were recruited to participate in a longitudinal study and completed measures of pain and pain impact at 5 time points (11- and 21 point NRS, 5- and 16 point VRS, 6- and 7 point FPS). All of the current study participants came from the same population		CrV CsV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
	<p>(CrV) Concurrent validity: There was a strong association among all measures, with the 21-point NRS showing the most consistently strong (all r's > 0.80) association with the other measures. However, even the weakest association ($r = 0.59$), found between the NRS-11 and the FPS-7, indicated a great deal of overlap between these 2 measures.</p> <p>(CsV) Factoranalyse: A single factor emerged. The loadings, all 0.90 or greater (except NRS-11 0.80), support the validity of each of the scales as measures of pain intensity.</p> <p>Convergent validity: Correlation coefficients between each of the 6 measures and measures of pain interference (a modified version of the Pain Interference Scale of the Brief Pain Inventory) and depression (Center for Epidemiological Studies–Depression Scale, CES-D), were all in the expected direction, although some variability in the coefficients can be seen. The 7-point Faces scale appeared to be most strongly associated with these 2 measures, the NRS-11 and VRS-5 showed the weakest associations with pain interference, and the VRS-5 showed the weakest association with depression. The difference between the association between the NRS-11 and pain interference ($r = 0.25$) and the 7-point Faces scale ($r = 0.50$) was statistically significant ($t(42) = 2.46$, $p < 0.05$, for the difference between coefficients).</p>	

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Kaasalainen, S. & Crook, J. (2003)	A 240-bed long-term-care facility in urban southwestern Ontario, Canada.	4 groups of 130 elderly long-term-care residents: (1) cognitively intact, (2) mildly cognitively impaired, (3) moderately cognitively impaired, and (4) extremely cognitively impaired. (n = 130)	Repeated measures design: FPS, NRS, Present Pain Intensity Scale (PPI) were conducted twice 48 hours apart.	S	CsV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Test-retest reliability: Test-retest reliability for the three verbal-report scales was moderate to strong for the cognitively intact group (FPS: ICC = 0.84; PPI: ICC = 0.55; NRS: ICC = 0.87) but decreased for the other groups. In addition, the error variances were low for the cognitively intact group (FPS: s² error = 0.53; PPI: s² error = 0.71; NRS: s² error = 1.45) but increased with increasing cognitive impairment.</p>	<p>(CsV) Convergent validity: The Pearson r correlations of the Pain Assessment in the communicatively Impaired (PACI) tool, a behavioural-observation measure, with the three verbal-report scales (FPS, PPI, NRS) were low to moderate. For the cognitively intact group, all of these correlations were moderate and significant (FPS: r = 0.66, p < 0.001; PPI: r = 0.62, p < 0.01; NRS: r = 0.65, p < 0.01). For the mildly impaired group, none were significant at the p < 0.05 level. For the moderately impaired group, the PACI correlated moderately and significantly with the FPS (r = 0.63, p < 0.001) and PPI (r = 0.64, p < 0.001). However, the correlation between the PACI and NRS for those with moderate impairment was low and non significant (r = 0.30, p < 0.12).</p>	<p>Test-retest for the 3 verbal-report scales was moderate to strong for elderly persons with no cognitive impairment but decreased for the other groups. Similarly, error variances were low for those with no cognitive impairment but increased with increasing cognitive impairment. These findings indicate that the level of cognitive impairment decreases the reliability of verbal reports of pain.</p> <p>Test-retest reliability for both the NRS and the FPS was strong for residents without cognitive impairment but declined considerably for those with mild and moderate impairment, suggesting that these tools may not be good choices for use with these two groups.</p> <p>For those with moderate cognitive impairment, the PPI seems to be a more appropriate and reliable tool than the FPS or the NRS.</p>

Betrouwbaarheid: Stability (S), Internal consistency (IC), Equivalence (E)

Validiteit: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Pautex, S., Michon, A., Guedira, M., Emond, H., Le Lous, P., Samaras, D. et al. (2006)	Geriatrics hospital and a geriatric psychiatry service.	129 patients aged 65 and older hospitalized during a 15 month-period who met DSM criteria for dementia, with a Mini-Mental State Examination score less than 11 and a Clinical Dementia Rating of 3 or greater. (n = 129)	Repeated measures design. Patients were asked to indicate their current level of pain on different scales (VAS, Faces Pain Scale and Verbal Rating Scale). This was repeated 30 minutes later either by the same investigator (50% of the cases) or by a different examiner who was blinded to the first assessment.	S E	
Paice, J. A. & Cohen, F. L. (1997)	A large tertiary care hospital.	A convenience sample of 50 adult oncological patients with pain. (n = 50)	Comparative study: the SDS was compared with VAS and NRS.		CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Intra-rater reliability: $R = 0.96$ ($p < 0.001$)</p> <p>(E) Interrater reliability: $R = 0.95$ ($p < 0.001$)</p>	<p>(CrV) Concurrent validity:</p> <ul style="list-style-type: none"> - Correlation between VRS and different scales at first assessment: <ul style="list-style-type: none"> 7. Correlation VRS – FPS: $r = 0.80$ ($p < 0.001$) 8. Correlation VRS – VAS: $r = 0.67$ ($p < 0.001$) 9. Correlation VRS – Doloplus 2 (observational pain scale): $r = 0.47$ ($p < 0.001$) - Correlation between VRS and different scales at second assessment: <ul style="list-style-type: none"> 10. Correlation VRS – FPS: $r = 0.79$ ($p < 0.001$) 11. Correlation VRS – VAS: $r = 0.73$ ($p < 0.001$) 12. Correlation VRS – Doloplus 2 (observational pain scale): $r = 0.63$ ($p < 0.001$) 	<p>A significant better comprehension of the Verbal Rating Scale (VRS) and the Faces Pain Scale (FPS) scale was found. This might suggest that the VAS is not the most appropriate pain assessment scale for patients with severe dementia.</p>
	<p>(CrV) Concurrent validity:</p> <ul style="list-style-type: none"> -The strong positive correlation between the SDS and the NRS was statistically significant: $r = 0.83$, $p < 0.001$. -SDS was also correlated to VAS: $r = 0.71$, $p < 0.001$ 	<p>A majority of subjects (50%) preferred the use of the NRS when comparing the three scales used to measure pain intensity. Fewer patients preferred the SDS (38%), with the VAS chosen least often (12%).</p> <p>Eleven subjects (20%) in this study were unable to complete the VAS or did so with great difficulty. All subjects were able to complete the NRS and SDS without apparent difficulty.</p>

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Herr, K. A., Spratt, K., Mobily, P. R., & Richardson, G. (2004)	Subjects were recruited through college bulletin board displays, community faith centers, senior-citizen centers, senior-housing, and long-term facilities.	86 younger adults (age 25-55) and 89 older adults (age 65-94). (n = 175)	Repeated measures design. Noxious heat stimuli were delivered to the ventral forearm by an electronically controlled contact thermode. The heat stimuli were programmed to last 5 seconds and to present randomly 43°C, 45°C, 46°C, 47°C, 48°C, 49°C or 51°C. A 2 minute trial interval followed each stimulus presentation during which the subject rated the stimulus on 5 scales.	IC E	CsV Sen

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(IC) Internal consistency: Intercorrelations between the scales were all statistically significant at every temperature ($p < 0.001$). <i>Cronbach's alpha within each scale across the 7 temperature values:</i> VAS: 0.88 NRS: 0.88 VDS: 0.86 VNS: 0.88 FPS: 0.88 <i>Cronbach's alpha within temperature across the 5 scales:</i> 43°C: 0.96 45°C: 0.96 46°C: 0.96 47°C: 0.97 48°C: 0.97 49°C: 0.97 51°C: 0.97 (E) Interrater reliability: VAS: 93.5% agreement NRS: 100% agreement VDS: 100% agreement VNS: 100% agreement FPS: 100% agreement</p>	<p>(CsV) Principal components analysis: A one-factor model was extracted from the different measurement scales. The correlation of each scale to the isolated factor was as follows: VAS: 0.94 NRS: 0.96 VDS: 0.95 VNS: 0.95 FPS: 0.86 (Sen) Each tool demonstrated significant increases in score associated with increase in temperature ($p < 0.001$).</p>	<p>The VAS had a significant higher failure rate (6.7%) in comparison with the NRS, VDS, VNS, and FPS.</p> <p>The VNS demonstrated significantly higher levels of pain report than the other 4 scales.</p> <p>The psychometric scale evaluation was conducted using an experimental pain stimulus instead of using clinical pain stimuli.</p> <p>The scales most preferred in order by the total sample: NRS (35.3%); VDS (25.3%); VNS (15.9%); FPS (12.9%); VAS (10.6%).</p>

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Taylor, L. J. & Herr, K. (2003)	Not specified.	A convenience sample of 57 volunteers age 58 and older. Seventy-seven percent (n = 44) of the sample scored 24 or less on the mental status exam, indicating some degree of cognitive impairment. The remaining 23% (n = 13) were cognitively intact. (n = 57)	Comparative study: Subjects were instructed to recall a vividly remembered pain and rate this remembered pain using the FPS, the VDS, the NRS and the Iowa Pain Thermometer (IPT).	S	CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Test-retest reliability: The researcher returned 2 weeks later and reminded subjects of the “vividly remembered pain” identified at the first assessment and asked them to rate that pain again using all four scales. Spearman rank correlation coefficients between the 2-week vividly remembered pain ratings ranged from 0.52 to 0.83 in both groups. In the cognitively impaired group, the FPS had the strongest reliability coefficient (0.79), followed by the VDS (0.63), NRS (0.57) and IPT (0.52). In the intact group, the strongest correlation was noted with the IPT (0.83), followed by the FPS (0.81), NRS (0.74), and VDS (0.73).</p>	<p>(CrV) Concurrent validity: Spearman correlations between present pain ratings on the selected scales were statistically significant ($p = 0.01$) and ranged from 0.81 to 0.96 in the intact group and from 0.74 to 0.83 in the impaired group. The lowest correlation was found between the FPS and VDS ($r = 0.74$).</p>	<p>All of the participants were able to use each of the pain intensity scales to rate their present pain in a manner that allowed interpretation of a single pain score (e.g., not selecting more than one response, selection of options outside the scale range). No failures were noted.</p> <p>The FPS showed strong test-retest stability in the cognitively impaired minority older adults.</p> <p>Of the 35 older adults who identified a scale preference, the FPS was the preferred scale in both the intact group ($n = 5, 62.5\%$) and the impaired group ($n = 15, 56\%$). However, it should be noted that almost 40% of both groups had no preference for a specific pain intensity scale.</p>

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Benaim, C., Froger, J., Cazottes, C., Gueben, D., Porte, M., Desnuelle, C., & Pelissier, J. Y. (2007)	2 rehabilitation units.	Patients who suffered a first unilateral middle cerebral artery stroke. A distinction was made between left and right hemispheric stroke patients (LHSP –RHSP). (n = 127)	5 year period prospective study. FPS was compared with vertical VAS and Verbal Ratings Scale.	S E	CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) Intra-rater reliability (n = 33): Kappa coefficients for FPS were 0.74 (0.13) and 0.53 (0.10) in LHSP and RHSP, respectively. Kappa coefficients for VRS were 0.39 (0.14) and 0.57 (0.15) in LHSP and in RHSP, respectively. ICC for VAS were 0.78 (0.46–0.92) and 0.90 (0.74–0.96) in LHSP and in RHSP, respectively.</p> <p>(E) Interrater reliability (n = 43): Kappa coefficients for FPS were 0.64 (standard error = 0.11) and 0.44 (0.09) in LHSP and RHSP, respectively. Kappa coefficients for VRS were 0.46 (0.12) and 0.52 (0.12) in LHSP and in RHSP, respectively. ICC for VAS were 0.72 (95% CI = 0.44–0.88) and 0.86 (0.68–0.94) in LHSP and in RHSP, respectively.</p>	<p>(CrV) Concurrent validity (n = 51): LHSP scores on the FPS were highly correlated with VAS ($r = 0.82$, $p < 0.001$) and with VRS scores ($r = 0.65$, $p < 0.01$). In the RHSP group, correlations were also high ($r = 0.72$, 0.72; $p < 0.001$ respectively).</p>	Among 71 patients, most LHSP preferred FPS (16/32) to VAS (6/32) and VRS (10/32), most RHSP preferred VAS (18/39) to FPS (11/39) and VRS (10/39). The difference was statistically significant ($p < 0.05$).

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Author (year)	Setting	Sample (n)	Design	Reliability	Validity
Pautex, S., Herrmann, F., Le Lous, P., Fabjan, M., Michel, J. P., & Gold, G. (2005)	The inpatient dementia consultation of the Geneva Geriatric Hospital.	Elderly who met DSM criteria for dementia. (n = 160)	Repeated measures design. Patients were asked to indicate their current level of pain on different scales (horizontal/ vertical VAS, Faces Pain Scale and Verbal Rating Scale). This was repeated 30 minutes later either by the same investigator (50% of the cases) or by a different examiner who was blinded to the first assessment.	S E	CrV

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Results reliability	Results validity	Commentary
<p>(S) <i>Intra-rater reliability</i> VRS: $r = 0.97$ ($p < 0.001$)</p> <p>(E) <i>Interrater reliability</i>: $r = 0.80$ ($p < 0.001$)</p>	<p>(CrV) <i>Concurrent validity</i>:</p> <ul style="list-style-type: none"> - Correlation between VRS and different scales at first assessment: <ol style="list-style-type: none"> 5. Correlation VRS – FPS: $r = 0.89$ ($p < 0.001$) 6. Correlation VRS – horizontal VAS: $r = 0.91$ ($p < 0.001$) 7. Correlation VRS – vertical VAS: $r = 0.89$ ($p < 0.001$) 8. Correlation VRS – Doloplus 2 (observational pain scale): $r = 0.31$ ($p < 0.001$) - Correlation between VRS and different scales at second assessment: <ol style="list-style-type: none"> 5. Correlation VRS – FPS: $r = 0.89$ ($p < 0.001$) 6. Correlation VRS – horizontal VAS: $r = 0.85$ ($p < 0.001$) 7. Correlation VRS – vertical VAS: $r = 0.89$ ($p < 0.001$) 8. Correlation VRS – Doloplus 2 (observational pain scale): $r = 0.34$ ($p < 0.001$) 	

Reliability: Stability (S), Internal consistency (IC), Equivalence (E)

Validity: Face validity (FV), Content validity (CtV), Criterion validity (CrV), Construct validity (CsV)

Sensitivity (Sen), Specificity (Sp), Positive Predictive Value (PPV), Negative Predictive Value (NPV), Receiver Operating Curve (ROC), Likelihood Ratio (LR), Odds Ratio (OR)

Verbal Rating Scale

Bron: Herr, K. A. & Mobily, P. R. (1993). Comparison of selected pain assessment tools for use with the elderly. *Appl.Nurs.Res.*, 6, 39-46.

Instructions: Have the patient circle the words to indicate the intensity of current pain. Document the corresponding number that the patient marks on this tool. A response of "No Pain" is given a value of zero, whereas a response of "Pain As Bad As It Could Be" is given a value of six. Evaluating the numerical value of the pain score or the descriptor selected by the patient over time helps determine whether pain intensity is less following pharmacological and/or nonpharmacological treatments.

Pain as bad as it could be

Extreme pain

Severe pain

Moderate pain

Mild pain

Slight pain

No pain

Reprinted with permission from the Dr. K. Herr.

Verbal Rating Scale

Instructies: Vraag de patiënt om het woord te omcirkelen dat de huidige pijnintensiteit weerspiegeld. Noteer het nummer dat overeenkomt met het antwoord dat de patiënt heeft aangeduid. Aan het antwoord "geen pijn" wordt een waarde van nul toegekend. Het antwoord "Ergste pijn die u zich kan voorstellen", krijgt een waarde gelijk aan zes. Indien men deze numerieke waarde of de door de patiënt gekozen omschrijving van de pijn gedurende een bepaalde periode evalueert, dan is dit een hulpmiddel bij het beoordelen van de pijnintensiteit na een al dan niet farmacologische behandeling.

Ergste pijn die u zich kan voorstellen

Extreme pijn

Ernstige pijn

Tamelijk veel pijn

Enige pijn

Nauwelijks pijn

Geen pijn

Wat is BEST?

BEST staat voor BElgian Screening Tools en is een studie uitgevoerd door de Universiteit Gent, afdeling Verplegingswetenschap in opdracht van de Federale Overheidsdienst Volksgezondheid, Veiligheid van de voedselketen en Leefmilieu, en in samenwerking met de Université Catholique de Louvain.

Doel van BeST

Bedoeling van het project is het oprichten van een databank met wetenschappelijk gevalideerde meetinstrumenten. Met het oog op het objectiveren van de diagnostiek en van de resultaten van verpleegkundige interventies, zijn valide en betrouwbare meetinstrumenten een basisvoorwaarde om effectieve verpleegkundige zorg te kunnen bieden. Onze aandacht gaat uit naar meetinstrumenten voor de verpleegkundige interventies die bij de Minimale Verpleegkundige Gegevens gescoord worden.

Wat kan u vinden in dit rapport?

In dit rapport wordt de inhoud van het project alsook de gehanteerde methodologie beschreven. Vervolgens worden de verschillende meetinstrumenten per thema besproken. Bovendien wordt het instrument ter beschikking gesteld indien we hertoe toestemming verkregen. Meetinstrumenten met een hoge betrouwbaarheid en validiteit werden tevens naar het Nederlands en het Frans vertaald.

Projectleiders UGent:

Prof. dr. T. Defloor
Prof. dr. M. Grypdonck

Projectmedewerkers UGent:

M. Daem
Dr. K. Vanderwee

Projectleider UCL:

Dr. M. Gobert

Projectmedewerkers UCL:

C. Piron

Projectleider FOD:

B. Folens

Projectmedewerkers FOD:

M. Lardennois

Gelieve bij elk gebruik van dit rapport als volgt te refereren:

Daem, M., Piron, C., Lardennois, M., Gobert, M., Folens, B., Vanderwee, K., Grypdonck, M., & Defloor T. (2007). Opzetten van een databank met gevalideerde meetinstrumenten: BEST-project. Brussel, Federale Overheidsdienst Volksgezondheid, Veiligheid van de voedselketen en Leefmilieu.